

Effects of Revascularisation on Evoked Cerebral Blood Oxygenation Responses in Stroke Patients

Shin Nakamura, Kaoru Sakatani, Tsuneo Kano, Tetsuya Hoshino, Norio Fujiwara, Yoshihiro Murata, and Yoichi Katayama

Abstract We demonstrated that ischemic strokes exhibit an increase of deoxyhemoglobin during activation. We evaluated the effect of revascularization on the abnormal evoked cerebral blood oxygenation (CBO) responses in these patients, employing near-infrared spectroscopy (NIRS). We selected five patients who exhibited an increase of deoxyhemoglobin associated with increases of oxyhemoglobin and total hemoglobin during activation for this study. These patients showed marked reductions of base-line regional cerebral blood flow and cerebrovascular reserve capacity, which were improved 1 week after revascularization. Postoperative NIRS demonstrated that the increase of deoxyhemoglobin during activation was not observed after revascularization. This preliminary study demonstrated that the abnormal evoked-CBO response in ischemic stroke patients could be improved by revascularization.

1 Introduction

Blood oxygenation level-dependent (BOLD) contrast functional MRI (BOLD-fMRI) has been used in functional studies on stroke patients, on the assumption that these patients have normal neurovascular coupling [1–3]. Recent studies have revealed, however, that BOLD-fMRI does not correctly image activation areas in stroke patients [4–10]. It was suggested that impairments of neurovascular coupling may alter the evoked cerebral blood oxygenation (CBO) responses and hemodynamic changes in stroke patients, and this could result in failure of BOLD imaging. BOLD-fMRI alone, however, cannot elucidate the precise mechanisms involved, since BOLD-fMRI provides information mainly about concentration changes of deoxyhemoglobin (deoxy-Hb), which is paramagnetic [11].

S. Nakamura (*)

Department of Neurosurgery, Nihon University School of Medicine, Tokyo, Japan
e-mail: nakashin51@ybb.ne.jp

Recently, we have evaluated evoked CBO changes in the primary sensorimotor cortex (PSMC) of patients with ischemic stroke [12, 13], employing near-infrared spectroscopy (NIRS) which allows measurements of concentration changes of not only deoxy-Hb, but also oxyhemoglobin (oxy-Hb) [14]. NIRS demonstrated a decrease of deoxy-Hb with increases of oxy-Hb and total hemoglobin (= sum of deoxy-Hb and oxy-Hb, t-Hb) in the activated PSMC on the non-lesion side, which is consistent with the physiological basis of BOLD imaging [11]. However, in the PSMC on the lesion side, the concentration of deoxy-Hb increased during the entire course of activation, concomitantly with increases of oxy-Hb and t-Hb. In addition, BOLD-fMRI showed significantly smaller activation volumes in the PSMC on the lesion side. These findings suggest that the increase of paramagnetic deoxy-Hb during activation caused a reduction of BOLD signal, and this resulted in the failure of BOLD imaging in stroke patients [14].

We hypothesized that the abnormal evoked CBO response was caused by decreases of baseline CBF and cerebrovascular response (CVR) in ischemic stroke patients. In order to test this hypothesis, we evaluated the effects of revascularization on the evoked CBO changes in ischemic strokes.

2 Methods

We studied 5 patients with cerebral ischemia (four males, one female, 65.6 ± 10.6 years). All of the patients had suffered hemodynamic compromise caused by occlusion of the internal carotid artery (ICA) in four cases and stenosis of the ICA in one case. Four patients received superficial temporal artery-middle cerebral artery (STA-MCA) anastomosis, while one patient received carotid endarterectomy (CEA). The patients did not show neurological deficits at the time of examination.

We evaluated the rCBF at rest and the %CVR response to acetazolamide (1.0 g; ACZ) in the territory of the MCA, employing single photon emission tomography (SPECT) (PRISM 2000XP, Shimadzu Co., Japan): $\%CVR = \{(rCBF_{ACZ} - rCBF_{rest}) / rCBF_{rest}\} \times 100$, where rCBF_{ACZ} and rCBF_{rest} represent the rCBF before and after the injection of ACZ, respectively.

We measured the evoked CBO responses in the PSMC contralateral to the task performance using a multi-channel NIRS (OMM 2000, Shimadzu Co., Japan). This system consists of 16 light-source fibers and 16 detectors, resulting in 48 source-detector pairs; each light source has three laser diodes with wavelengths of 780, 805, and 830 nm [15, 16]. The optodes for the NIRS topography were placed on the skull to cover the motor cortex, employing a holder cap to avoid motion-related artifacts; the distance between each optode was 30 mm. We measured the concentration changes in oxy-Hb, deoxy-Hb, and t-Hb in the PSMC on the lesion side during contralateral hand-grasping tasks. The task paradigm consisted of 40 s of rest and 40 s of self-paced hand grasping; this task-rest cycle was repeated 6 times.

We evaluated the changes in oxy-Hb, deoxy-Hb, and t-Hb by subtracting the mean baseline values (40 s) from the mean stimulation values (40 s). Comparisons were made for each of the NIRS parameters using paired *t*-tests ($p < 0.05$ was defined as the criterion of a significant difference).

We classified NIRS responses into three patterns depending on concentration changes of deoxy-Hb during activation. That is, the deoxy-Hb concentration significantly decreased (Pattern 1), did not change (Pattern 2), or significantly increased (Pattern 3). Both oxy-Hb and t-Hb increased during the task in all subjects.

3 Results

Figure 1 shows an example of evoked CBO changes before and after revascularization. Note that revascularization reversed the direction of deoxy-Hb changes, and increased the extent of the increases in oxy-Hb and t-Hb. Pre-operative NIRS demonstrated an increase of deoxy-Hb associated with increases of oxy-Hb and t-Hb (Pattern 3) during activation in all patients,




Fig. 1 NIRS topographic maps of changes in oxy-Hb and deoxy-Hb during right grasping task overlaid on anatomical MRI surface images before (a) and after (b) revascularization. The circles indicate the ROI for analysis of NIRS parameter changes (right). Note that revascularization reversed the direction of deoxy-Hb changes, and increased the extent of the increases in oxy-Hb and t-Hb

while postoperative NIRS demonstrated a decrease of deoxy-Hb (Pattern 1) in three patients and no change of deoxy-Hb (Pattern 2) in two subjects.

abnormal evoked CBO response required only 1 week after revascularization, suggesting that the abnormal evoked CBO response was due to hemodynamic effects, rather than structural changes of cerebral vessels caused by arteriosclerosis.

References

1. Cramer SC, Nelles G, Benson RR et al. (1997) A functional MRI study of subjects recovered from hemiparetic stroke. *Stroke* 28:2518–2527
2. Cao Y, D’Olhaberriague L, Vikingstad EM et al. (1998) Pilot study of functional MRI to assess cerebral activation of motor function after poststroke hemiparesis. *Stroke* 29:112–122
3. Kato H, Izumiyama M, Koizumi H et al. (2002) Near-infrared spectroscopic topography as a tool to monitor motor reorganization after hemiparetic stroke: A comparison with functional MRI. *Stroke* 33:2032–2036
4. Pineiro R, Pendlebury S, Johansen-Berg H et al. (2002) Altered hemodynamic responses in patients after subcortical stroke measured by functional MRI. *Stroke* 33:103–109
5. Røther J, Knab R, Hamzei F et al. (2002) Negative dip in BOLD fMRI is caused by blood flow-oxygen consumption uncoupling in humans. *NeuroImage* 15:98–102
6. D’Esposito M, Deouell LY, Gazzaley A (2003) Alterations in the BOLD fMRI signal with aging and disease: A challenge for neuroimaging. *Nature Rev Neurosci* 4:863–872
7. Hamzei F, Knab R, Weiller C et al. (2003) The influence of extra- and intracranial artery disease on the BOLD signal in FMRI. *NeuroImage* 20:1393–1399
8. Rossini PM, Altamura C, Ferretti A et al. (2004) Does cerebrovascular disease affect the coupling between neuronal activity and local haemodynamics? *Brain* 127:99–110
9. Binkofski F, Seitz RJ (2004) Modulation of the BOLD-response in early recovery from sensorimotor stroke. *Neurology* 63:1223–1229
10. Krainik A, Hund-Georgiadis M, Zysset S et al. (2005) Regional impairment of cerebrovascular reactivity and BOLD signal in adults after stroke. *Stroke* 36:1146–1152
11. Ogawa S, Lee TM, Kay AR et al. (1990) Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc Natl Acad Sci USA* 87:9868–9872
12. Murata Y, Sakatani K, Hoshino T et al. (2006) Effects of cerebral ischemia on evoked cerebral blood oxygenation responses and BOLD contrast functional MRI in stroke patients. *Stroke* 37:2514–2520
13. Sakatani K, Murata Y, Fujiwara N et al. (2007) Comparison of blood-oxygen-level-dependent functional magnetic resonance imaging and near-infrared spectroscopy recording during functional brain activation in patients with stroke and brain tumors. *J Biomed Optics* 12:062110
14. Jöbsis FF (1977) Noninvasive, infrared monitoring of cerebral and myocardial oxygen sufficiency and circulatory parameters. *Science* 198:1264–1267
15. Hoshino T, Sakatani K, Katayama Y et al. (2005) Application of multi-channel near infrared spectroscopic topography to physiological monitoring of the cortex during cortical mapping: Technical case report. *Surg Neurol* 64:272–275
16. Hoshino T, Katayama Y, Sakatani K et al. (2006) Intraoperative monitoring of cerebral blood oxygenation and hemodynamics during EC-IC bypass surgery by a newly developed visual light spectroscopy system. *Surg Neurol* 65:569–576
17. Anderson DE, McLane MP, Reichman OH et al. (1992) Improved cerebral blood flow and CO₂ reactivity after microvascular anastomosis in patients at high risk for recurrent stroke. *Neurosurgery* 31:26–33
18. Gibbs JM, Wise RJ, Thomas DJ et al. (1987) Cerebral hemodynamic changes after extracranial-intracranial bypass surgery. *J Neurol Neurosurg Psychiatry* 50:140–150

19. Powers WJ, Martin WR, Herscovitch P et al. (1984) Extracranial-intracranial bypass surgery: hemodynamic and metabolic effects. *Neurology* 34:1168–1174
20. Boxerman JI, Bandettini PA, Kwong KK et al. (1995) The intravascular contribution to fMRI signal change: Monte Carlo modeling and diffusion-weighted studies in vivo. *Magn Reson Med* 34:4–10